



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K</b>	<b>A2</b>	(11) International Publication Number: <b>WO 00/51546</b> (43) International Publication Date: 8 September 2000 (08.09.00)									
<p>(21) International Application Number: PCT/US00/05109</p> <p>(22) International Filing Date: 29 February 2000 (29.02.00)</p> <p>(30) Priority Data:</p> <table border="0"><tr><td>60/122,277</td><td>1 March 1999 (01.03.99)</td><td>US</td></tr><tr><td>60/148,324</td><td>11 August 1999 (11.08.99)</td><td>US</td></tr><tr><td>Not furnished</td><td>22 February 2000 (22.02.00)</td><td>US</td></tr></table> <p>(71) Applicant: SEPRACOR INC. [US/US]; 111 Locke Drive, Marlborough, MA 01752 (US).</p> <p>(72) Inventors: JERUSSI, Thomas, P.; 19 Garvey Road, Framingham, MA 01701 (US). MCCULLOUGH, John, R.; 6 Davidson Road, Worcester, MA 01605 (US). SENANAYAKE, Chisantha, H.; 11 Old Farm Circle, Shrewsbury, MA 01545 (US). FANG, Qun, K.; 35 Atwood Street, Wellesley, MA 02181 (US).</p> <p>(74) Agents: INSOGNA, Anthony, M. et al.; Pennie &amp; Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).</p>		60/122,277	1 March 1999 (01.03.99)	US	60/148,324	11 August 1999 (11.08.99)	US	Not furnished	22 February 2000 (22.02.00)	US	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published</p> <p><i>Without international search report and to be republished upon receipt of that report.</i></p>
60/122,277	1 March 1999 (01.03.99)	US									
60/148,324	11 August 1999 (11.08.99)	US									
Not furnished	22 February 2000 (22.02.00)	US									
<p>(54) Title: BUPROPION METABOLITES AND METHODS OF THEIR SYNTHESIS AND USE</p> <p>(57) Abstract</p> <p>Methods and compositions are disclosed which utilize metabolites of bupropion for treating disorders ameliorated by inhibition of neuronal monoamine reuptake. Such disorders include, but are not limited to, erectile dysfunction, affective disorders, cerebral function disorders, cigarette smoking, and incontinence. The invention further discloses methods of making optically pure bupropion metabolites.</p>											

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

**BUPROPION METABOLITES AND METHODS  
OF THEIR SYNTHESIS AND USE**

**1. FIELD OF THE INVENTION**


5           This invention relates to synthesis of, methods of using, and compositions comprising bupropion metabolites and isomers thereof.

**2. BACKGROUND OF THE INVENTION**

10           Bupropion, a racemic mixture of (+)- and (-)-1-(3-chlorophenyl)-2-[(1,1-dimethylethyl)amino]-1-propanone, is an antidepressant of the aminoketone class, and is described in U.S. Patent Nos. 3,819,706 and 3,885,046. The hydrochloride salt of bupropion is sold under the trade names WELLBUTRIN® and WELLBUTRIN SR® for the treatment of depression. Bupropion is also sold under the trade name ZYBAN® as a drug useful to achieve smoking cessation. Additional benefits of bupropion maleate are reported  
15           in European Patent Application No. 118036.


          Although its mechanism of action is poorly understood, bupropion is reportedly a weak but selective inhibitor of dopamine. Its potency as an inhibitor of norepinephrine reuptake is reportedly only half of that for dopamine, and it shows little affinity for the serotonergic transport system. Ascher, J. A., *et al.*, *J. Clin. Psychiatry*, 56:395-401 (1995).

20           Bupropion is extensively metabolized in man and animal. Three metabolites found in the plasma of healthy humans to which it has been administered are shown in Scheme 1:



Posner, J., *et al.*, *Eur. J. Clin Pharmacol.*, 29:97-103 (1985); Suckow, R.F., *et al.*, *Biomedical Chromatography*, 11:174-179 (1997). Referring to Scheme 1, metabolite 1 has the chemical name 2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; metabolite 2 has the chemical name 1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; and  
 5 metabolite 3 has the chemical name 1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanone. Because bupropion is racemic and its metabolites are chiral, enantiomers of each of the metabolites 1, 2, and 3 likely exist in human plasma following its administration.

The bupropion metabolite 1, often referred to as "hydroxybupropion," has two chiral  
 10 carbon atoms and can thus exist as two pairs of enantiomers. These are shown in Scheme 2:



Pair 1a is reportedly the most active human metabolite bupropion. Kelley, J. L., *et al.*, *J. Med. Chem.*, 39:347-349 (1996). The mixture 1a has been isolated from human plasma and allegedly separated into its (S,S) and (R,R) components. Suckow, R.F., *et al.*, *Biomedical Chromatography*, 11:174-179 (1997).  
 20

The amino alcohol metabolite 2 can also exist as two pairs of enantiomers. The pair wherein the alcohol and amine moieties are *cis* each other is commonly referred to as the  
 25 *erythro*-amino alcohol metabolite; the pair wherein the two moieties are *trans* to each other is referred to as the *threo*-amino alcohol metabolite.

The *tert*-butyl alcohol metabolite 3 can exist as one of two enantiomers. This metabolite, the accumulation of which in human plasma coincides with the elimination of a single dose of bupropion, is believed by some to be a precursor to hydroxybupropion.  
 30 Posner, J., *et al.*, *Eur. J. Clin. Pharmacol.*, 29:97-103 (1985); Suckow, R.F., *et al.*, *Biomedical Chromatography*, 11:174-179 (1997).

Clearly, the metabolism of bupropion, which is complicated and poorly understood, results in a complex array of optically active compounds. The structures of these molecules and their chirality provides the skilled artisan with difficult issues of asymmetric synthesis, chiral resolution, and pharmacological activity.

5           Bupropion is widely used to treat affective disorders in patients who do not respond to, or cannot tolerate, other antidepressants such as tricyclic agents or monoamine oxidase inhibitors. Examples of affective disorders are depression and bipolar manic-depression. Bupropion is also useful in the treatment of other diseases or conditions associated with the reuptake of neuronal monoamines such as serotonin and norepinephrine. These reportedly  
10   include: schizophrenia (U.S. Patent No. 5,447,948); attention-deficit disorder; psychosexual dysfunction (U.S. Patent No. 4,507,323); bulimia and other eating disorders; Parkinson's disease; migraine (U.S. Patent No. 5,753,712); and chronic pain. Bupropion also reportedly increases success rates in some smoking cessation treatments. Rose, J.E., *Annu. Rev. Med.*, 47:493-507 (1996); Ferry, L.H. *et al.*, *J. Addict. Dis.*, 13:A9 (1994); and  
15   Lief, H.I., *Am. J. Psychiatry*, 153(3):442 (1996).

          Further uses of bupropion reportedly include the treatment of: the effects of ethanol (U.S. Patent No. 4,393,078); tardive dyskinesia (U.S. Patent No. 4,425,363); drowsiness (U.S. Patent Nos. 4,571,395 and 4,798,826); minimal brain dysfunction (U.S. Patent No. 4,435,449); psychosexual dysfunction (U.S. Patent No. 4,507,323); prostate hypertrophy  
20   and sexual dysfunction (U.S. Patent No. 4,835,147); psychostimulant addiction (U.S. Patent No. 4,935,429); substance abuse (U.S. Patent No. 5,217,987); high cholesterol (U.S. Patent No. 4,438,138); and weight gain (U.S. Patent No. 4,895,845).

          Certain advantages exist in using bupropion for the treatment of diseases and conditions such as those provided above. For example, it does not inhibit monoamine  
25   oxidase or block the reuptake of serotonin, unlike other neuronal monoamine reuptake inhibitors. Administration of bupropion can thus avoid or lessen many of the adverse side effects commonly associated with other antidepressants such as tricyclic agents and monoamine oxidase inhibitors.

          Unfortunately, bupropion is not free of adverse effects. Administration of the drug  
30   can cause seizures, especially in patients currently taking the monoamine oxidase inhibitor phenelzine. Other frequently reported adverse effects associated with the use of bupropion include nausea, vomiting, excitement, agitation, blurred or blurry vision, restlessness,

postural tremor, hallucinations/confusional states with the potential for abuse, anxiety, insomnia, headaches and/or migraines, dry mouth, constipation, tremor, sleeping disturbances, dermatologic problems (e.g., rashes), neuropsychiatric signs and symptoms (e.g., delusions and paranoia), and weight loss or gain. *Physicians' Desk Reference*® 1252-1258 (53<sup>rd</sup> ed. 1999). These effects are dose limiting in a number of patients, and can be particularly dangerous for Parkinson's patients.

There thus remains a need for a drug that provides the advantages of bupropion, but with fewer disadvantages. Compounds and pharmaceutical compositions are desired that can be used for the treatment and prevention of disorders and conditions while incurring fewer of the adverse effects associated with bupropion.

### 3. SUMMARY OF THE INVENTION

This invention encompasses methods of making and using bupropion metabolites and pharmaceutical compositions and dosage forms comprising bupropion metabolites. In particular, the invention provides methods of synthesizing optically pure (S,S)-hydroxybupropion and optically pure (R,R)-hydroxybupropion.

The invention further provides methods of treating and preventing conditions that include, but are not limited to, erectile dysfunction, affective disorders, cerebral function disorders, tobacco smoking, and incontinence. Methods of the invention comprise administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite, or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferred methods of the invention further comprise the use of at least one additional physiologically active agent such as a selective serotonin reuptake inhibitor ("SSRI"), 5-HT<sub>3</sub> antagonist, or nicotine with a bupropion metabolite.

Pharmaceutical compositions and dosage forms of the invention comprise a therapeutically or prophylactically effective amount of a bupropion metabolite and optionally at least one additional physiologically active agent such as a SSRI, 5-HT<sub>3</sub> antagonist, or nicotine.

#### 3.2. DEFINITIONS

As used herein, the term "patient" includes mammal, which includes human.

As used herein, the term "bupropion metabolite" includes, but is not limited to, 2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol, 1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol, and 1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanone. As used herein, the term "optically pure bupropion metabolite" includes, but is not limited to, optically pure: (R,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (S,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (R,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (R,R)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (S,R)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (S,S)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (R,S)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (R)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanone; and (S)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanone.

As used herein to describe a composition, the terms "substantially optically pure," "optically pure," and "optically pure enantiomer" mean that the composition contains greater than about 90% of the desired stereoisomer by weight, preferably greater than about 95% of the desired stereoisomer by weight, and most preferably greater than about 99% of the desired stereoisomer by weight, said weight percent based upon the total weight of bupropion metabolite. As used herein to describe a composition, the term "substantially free" means that the composition contains less than about 10% by weight, preferably less than about 5% by weight, and more preferably less than about 1% by weight of the undesired stereoisomer.

As used herein, the term "pharmaceutically acceptable salt" refers to a salt prepared from a pharmaceutically acceptable non-toxic inorganic or organic acid or base. The compounds of the invention that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. Acids that can be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds of the invention are those that form non-toxic acid addition salts, *i.e.*, salts containing pharmacologically acceptable anions, such as, but not limited to, hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, formate, acetate, propionate, succinate, camphorsulfonate, citrate, acid citrate, fumarate, gluconate, isethionate, lactate, malate,

mucate, gentisate, isonicotinate, saccharate, tartrate, bitartrate, para-toluenesulfonate, glycolate, glucuronate, maleate, furoate, glutamate, ascorbate, benzoate, anthranilate, salicylate, phenylacetate, mandelate, embonate (pamoate), methanesulfonate, ethanesulfonate, pantothenate, benzenesulfonate, stearate, sulfanilate, alginate, p-toluenesulfonate, and galacturonate. Particularly preferred anions are hydrobromide, hydrochloride, phosphate, acid phosphate, maleate, sulfate, and acid phosphate. Most particularly preferred anions are hydrochloride and maleate.

Compounds of the invention that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds of the invention are those that form non-toxic base addition salts, *i.e.*, salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), lysine, and procaine.

As used herein, the terms "avoiding adverse side effects" and "avoiding adverse effects" mean eliminating or reducing at least one adverse effect associated with the administration of a particular compound or mixture of compounds.

As used herein, the term "adverse side effects associated with racemic bupropion" includes, but is not limited to, seizures, nausea, vomiting, excitement, agitation, blurred or blurry vision, restlessness, postural tremor, hallucinations/confusional states with the potential for abuse, anxiety, insomnia, headaches and/or migraines, dry mouth, constipation, tremor, sleeping disturbances, dermatologic problems (*e.g.*, rashes), neuropsychiatric signs and symptoms (*e.g.*, delusions and paranoia), and weight gain.

As used herein, the term "adverse side effects associated with the inhibition of dopamine reuptake" includes, but is not limited to, seizures, nausea, vomiting, excitement, agitation, blurred or blurry vision, restlessness, postural tremor, hallucinations/confusional states with the potential for abuse, anxiety, insomnia, headaches and/or migraines, dry mouth, constipation, tremor, sleeping disturbances, dermatologic problems (*e.g.*, rashes), neuropsychiatric signs and symptoms (*e.g.*, delusions and paranoia), and weight gain.

As used herein, the term "disorder ameliorated by the inhibition of neuronal monoamine reuptake" and "disorder related to reuptake of neuronal monamines" mean an



acute or chronic disease, disorder, or condition having symptoms that are reduced or alleviated by the inhibition of neuronal monoamine reuptake, and especially by the inhibition of norepinephrine (or noradrenaline) and serotonin reuptake. Disorders ameliorated by inhibition of neuronal monoamine reuptake include, but are not limited to, 5 erectile dysfunction, affective disorders, cerebral function disorders, tobacco smoking, and incontinence.

As used herein, the term "affective disorder" includes, but is not limited to, depression, anxiety disorders, attention deficit disorder, attention deficit disorder with hyperactivity, bipolar and manic conditions, bulimia, obesity or weight-gain, narcolepsy, 10 chronic fatigue syndrome, seasonal affective disorder, premenstrual syndrome, substance addiction or abuse, and nicotine addiction.

As used herein, the term "substance addiction" includes, but is not limited to, addiction to cocaine, heroin, nicotine, alcohol, opioids, anxiolytic and hypnotic drugs, cannabis (marijuana), amphetamines, hallucinogens, phencyclidine, volatile solvents, and 15 volatile nitrites. Nicotine addiction includes nicotine addiction of all known forms, such as smoking cigarettes, cigars and/or pipes, and addiction to chewing tobacco.

As used herein, the terms "attention deficit disorder" (ADD), "attention deficit disorder with hyperactivity" (ADHD), and "attention deficit/hyperactivity disorder" (AD/HD), are used in accordance with their accepted meanings in the art. *See, e.g.,* 20 *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Ed., American Psychiatric Association, 1997 (DSM-IV™) and *Diagnostic and Statistical Manual of Mental Disorders*, 3<sup>rd</sup> Ed., American Psychiatric Association (1981) (DSM-III™).

As used herein, the term "depression" includes a disease or condition characterized by changes in mood, feelings of intense sadness, despair, mental slowing, loss of 25 concentration, pessimistic worry, agitation, and self-deprecation. Physical symptoms of depression that may be reduced or alleviated by the methods of the invention include insomnia, anorexia, weight loss, decreased energy and libido, and abnormal hormonal circadian rhythms.

As used herein, the term "cerebral function disorder" includes, but is not limited to, 30 cerebral function disorders involving intellectual deficits such as senile dementia, Alzheimer's type dementia, memory loss, amnesia/amnestic syndrome, epilepsy, disturbances of consciousness, coma, lowering of attention, speech disorders, Parkinson's

disease, Lennox syndrome, autism, hyperkinetic syndrome and schizophrenia. Also within the meaning of the term are disorders caused by cerebrovascular diseases including, but not limited to, cerebral infarction, cerebral bleeding, cerebral arteriosclerosis, cerebral venous thrombosis, head injuries, and the like where symptoms include disturbance of  
5 consciousness, senile dementia, coma, lowering of attention, and speech disorders.

As used herein, the term "method of treating Parkinson's disease" means relief from the symptoms of Parkinson's disease which include, but are not limited to, slowly increasing disability in purposeful movement, tremors, bradykinesia, rigidity, and a disturbance of posture.

10 As used herein, the term "a method for treating obesity or weight gain" means reduction of weight, relief from being overweight, relief from gaining weight, or relief from obesity, all of which are usually due to extensive consumption of food.

As used herein, the term "a method of treating or preventing incontinence" means prevention of or relief from the symptoms of incontinence including involuntary voiding of  
15 feces or urine, and dribbling or leakage of feces or urine which may be due to one or more causes including, but not limited to, pathology altering sphincter control, loss of cognitive function, overdistention of the bladder, hyper-reflexia and/or involuntary urethral relaxation, weakness of the muscles associated with the bladder, or neurologic abnormalities. As used herein, the term "urinary incontinence" encompasses stress urinary  
20 incontinence and urge urinary incontinence.

#### 4. DETAILED DESCRIPTION OF THE INVENTION

This invention relates to methods and compositions that inhibit the reuptake of neuronal monoamines (*e.g.*, norepinephrine). The invention thereby provides methods,  
25 pharmaceutical compositions, and dosage forms for the treatment or prevention of disorders that are ameliorated by the inhibition of neuronal monoamine reuptake including, but are not limited to, erectile dysfunction, affective disorders, cerebral function disorders, tobacco smoking, and incontinence.

The methods, pharmaceutical compositions, and dosage forms of the invention  
30 comprise a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferably the bupropion metabolite is optically pure. More preferably, the

bupropion metabolite is optically pure (S,S)-hydroxybupropion (*i.e.*, (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol).

5 The bupropion metabolite (S,S)-hydroxybupropion is an unexpectedly selective norepinephrine reuptake inhibitor that does not significantly inhibit dopamine reuptake. It can thus be used to treat or prevent disorders related to norepinephrine reuptake without incurring adverse side effects associated with the inhibition of dopamine reuptake. It can also be used to treat or prevent disorders related to norepinephrine reuptake while reducing or avoiding adverse effects associated with racemic bupropion.

10 A first embodiment of the invention is a method of treating or preventing a disorder that is ameliorated by the inhibition of neuronal monoamine reuptake which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite, or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferably, the bupropion metabolite is optically pure. More preferably, the bupropion metabolite is optically pure (S,S)-  
15 hydroxybupropion. In a preferred method encompassed by this embodiment, adverse effects associated with the inhibition of dopamine reuptake are reduced or avoided.

In another preferred method encompassed by this embodiment, the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively administered with an additional pharmacologically active compound, *i.e.*, the bupropion  
20 metabolite and an additional pharmacologically active compound are administered as a combination, concurrently but separately, or sequentially by any suitable route (*e.g.*, orally, transdermally, or mucosally).

Additional pharmacologically active compounds include, but are not limited to, SSRIs, 5-HT<sub>3</sub> inhibitors, and nicotine. Selective serotonin reuptake inhibitors are  
25 compounds that inhibit the central nervous system uptake of serotonin while having reduced or limited affinity for other neurologically active receptors. Examples of SSRIs include, but are not limited to, citalopram (CELEXA<sup>®</sup>); fluoxetine (PROZAC<sup>®</sup>) fluvoxamine (LUVOX<sup>®</sup>); paroxetine (PAXIL<sup>®</sup>); sertraline (ZOLOFT<sup>®</sup>); venlafaxine (EFFEXOR<sup>®</sup>); and optically pure stereoisomers, active metabolites, and pharmaceutically acceptable salts,  
30 solvates, and clathrates thereof.

Preferred 5-HT<sub>3</sub> antagonists are antiemetic agents. Examples of suitable 5-HT<sub>3</sub> antagonists include, but are not limited to, granisetron (KYTRIL<sup>®</sup>), metoclopramide

(REGLAN®), ondansetron (ZOFTRAN®), renzapride, zacopride, tropisetron, and optically pure stereoisomers, active metabolites, and pharmaceutically acceptable salts, solvates, and clathrates thereof.

5 A second embodiment of the invention encompasses a method of treating or preventing erectile dysfunction which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferably, the bupropion metabolite is optically pure. More preferably, the bupropion metabolite is optically pure (S,S)-hydroxybupropion.

10 In a preferred method encompassed by this embodiment, the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered transdermally or mucosally (e.g., nasally, sublingually, or buccally).

In another preferred method encompassed by this embodiment, the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively  
15 administered with a 5-HT<sub>3</sub> antagonist.

A third embodiment of the invention encompasses a method of treating or preventing an affective disorder which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.  
20 Preferably, the bupropion metabolite is optically pure. More preferably, the bupropion metabolite is optically pure (S,S)-hydroxybupropion.

A particular preferred method encompassed by this embodiment is a method of treating or preventing depression. Another preferred method encompassed by this embodiment is a method of treating or preventing narcolepsy. Yet another preferred  
25 method encompassed by this embodiment is a method of treating or preventing nicotine addiction.

A fourth embodiment of the invention encompasses a method of treating or preventing a cerebral function disorder which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a  
30 bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferably, the bupropion metabolite is optically pure. More preferably, the bupropion metabolite is optically pure (S,S)-hydroxybupropion.

A particular preferred method encompassed by this embodiment is a method of treating or preventing Parkinson's disease. Another preferred method encompassed by this embodiment is a method of treating or preventing epilepsy.

5 A fifth embodiment of the invention encompasses a method of eliciting smoking cessation which comprises administering to a patient who smokes tobacco a therapeutically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferably, the bupropion metabolite is optically pure. More preferably, the bupropion metabolite is optically pure (S,S)-hydroxybupropion.

10 In preferred method encompassed by this embodiment, the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered orally, mucosally, or transdermally. In a more preferred method, the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered transdermally.

15 In another preferred method encompassed by this embodiment, the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively administered with a therapeutically effective amount of nicotine. Preferably, the nicotine and/or bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered orally, mucosally, or transdermally. More preferably, the nicotine and/or bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered transdermally.

20 A sixth embodiment of the invention encompasses a method of treating or preventing incontinence which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferably, the bupropion metabolite is optically pure. More preferably, the bupropion  
25 metabolite is optically pure (S,S)-hydroxybupropion. A preferred method encompassed by this embodiment is a method of treating or preventing stress urinary incontinence. In another preferred method encompassed by this embodiment, the patient is a human of an age greater than 50 years or less than 13 years.

30 A seventh embodiment of the invention encompasses pharmaceutical compositions and dosage forms which comprise a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof. Preferably, the bupropion metabolite is optically pure. More preferably, the bupropion metabolite is optically pure (S,S)-hydroxybupropion.

Pharmaceutical compositions and dosage forms encompassed by this embodiment can further comprise at least one additional pharmacologically active compound.

Additional pharmacologically active compounds include, but are not limited to, SSRIs, 5-HT<sub>3</sub> inhibitors, and nicotine as described above.

5        An eighth embodiment of the invention encompasses a process for preparing optically pure (S,S)-hydroxybupropion which comprises: the asymmetric dihydroxylation of Z-1-(3-chlorophenyl)-1-*tert*-butyldimethylsilyloxy-1-propene to form an intermediate; the reaction of the intermediate with 2-amino-2-methyl-1-propanol to form (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; and the isolation of the (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol. Preferably, the intermediate formed  
10        by the asymmetric dihydroxylation is an  $\alpha$ -hydroxy ketone activated by trifluoromethane sulfonic anhydride.

A ninth embodiment of the invention encompasses a process for preparing optically pure (R,R)-hydroxybupropion which comprises: the asymmetric dihydroxylation of Z-1-(3-chlorophenyl)-1-*tert*-butyldimethylsilyloxy-1-propene to form an intermediate; the reaction  
15        of the intermediate with 2-amino-2-methyl-1-propanol to form (R,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; and the isolation of the (R,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol. Preferably, the intermediate formed by the asymmetric dihydroxylation is an  $\alpha$ -hydroxy ketone activated by trifluoromethane sulfonic  
20        anhydride.

#### 4.1 SYNTHESIS OF BUPROPION METABOLITES


The metabolism of bupropion, which varies among species, is complex and poorly understood. Bupropion has been shown to induce its own metabolism in mice, rats, and  
25        dogs, and may do so in human patients to whom the drug has been administered over long periods of time. In the plasma of healthy humans to which the drug has been administered, however, at least three major metabolites are found. *Physicians' Desk Reference*<sup>®</sup> 1252-1258 (53<sup>rd</sup> ed. 1999). Each of these major metabolites is chiral, meaning that a total of at least ten optically pure bupropion metabolites exist in varying concentrations in the plasma  
30        of a patient following administration of the drug.

It is possible to prepare a mixture of the stereoisomers of the amino alcohol metabolite of bupropion (*i.e.*, 1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-

propanol) using techniques known to those skilled in the art. *See, e.g.*, Japanese Patent No. 63091352. The optically pure forms of this metabolite can be isolated from the resulting mixture by any method known to those skilled in the art, including high performance liquid chromatography (HPLC) and the formation and crystallization of chiral salts. *See, e.g.*,  
5 Jacques, J., *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., *et al.*, *Tetrahedron*, 33:2725 (1977); Eliel, E. L., *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S. H., *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN, 1972).

10 It is also possible to prepare a mixture of the stereoisomers of the *tert*-butyl alcohol metabolite of bupropion (*i.e.*, 1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanone). From the resulting mixture of compounds, individual stereoisomers may be resolved using conventional means such as HPLC and the formation and crystallization of chiral salts.

15 Alleged analogues of the hydroxybupropion metabolite (*i.e.*, 2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol) have reportedly been prepared according to methods known to those skilled in the art. *See, e.g.*, Kelly, J. L., *et al.*, *J. Med. Chem.*, 39:347-349 (1996). Not until now, however, has an effective and efficient synthetic process been discovered for the synthesis of optically pure stereoisomers of the metabolite itself. This  
20 process utilizes a protected alcohol derivative of 1-(3-chlorophenyl)-1-propene, which is dihydroxylated and then cyclized to form the morpholinol moiety. A particular embodiment of this process which can be used to form optically pure (R,R)- and (S,S)-hydroxybupropion is shown in Scheme 3:



Scheme 3

According to a preferred embodiment of this process, compound 5 is prepared in step (a), wherein the ketone 4 is converted to its enolate, preferably by use of a strong base such as, but not limited to, lithium hexamethyldisilazide (LHMDS) and lithium diisopropylamide (LDA). A preferred base is LDA. The enolate is then trapped using a  
5 protecting agent such as, but not limited to, *tert*-butyl-dimethylsilyl chloride (TBSCl). Compound 5 is preferably isolated prior to step (b).

According to step (b), the vinyl group of compound 5 is asymmetrically dihydroxylated to give the ketone. It has been found that the choice of reagent used to asymmetrically hydroxylate compound 5 affects the stereochemistry of the resulting  
10 product, as well as its optical purity. Suitable asymmetric hydroxylation reagents include, for example, oxides of transition metals such as manganese and osmium, although preferred reagents are AD-mix- $\alpha$  and AD-mix- $\beta$ . These reagents have been found to selectively dihydroxylate the vinyl group of compound 5 to reform the ketone. Use of AD-mix- $\alpha$  yields (R)-3-chloro-2-hydroxyl-propiophenone (i), while use of AD-mix- $\beta$  yields (S)-3-  
15 chloro-2-hydroxyl-propiophenone. Although not necessary, it has been found that care taken to ensure the optical purity of the intermediate (*e.g.*, compound 6) formed in this step improves the optical purity of the final product (*i.e.*, optically pure hydroxybupropion). It is thus preferred that step (b) further include purification by, for example, column chromatography.

20 Substantially optically pure (5,5)-hydroxybupropion 7 is formed in step (c) of Scheme 3, which comprises the stereospecific displacement of triflates of compound 6:



wherein R is triflate (*i.e.*, -OSO<sub>2</sub>CF<sub>3</sub>). Other compounds potentially useful in the synthesis of the compounds of the invention are those wherein R is mesylate, tosylate, or nosylate.  
30 Substantially optically pure (R,R)-hydroxybupropion is preferably formed from the triflate of opposite stereochemistry.



Triflation is conducted with pyridine base. A preferred base is lutidine when used in combination with trifluoromethanesulfonic anhydride. The cyclized product **7** is isolated by extraction, and purified by chromatography. Substantially optically pure (R,R)-hydroxybupropion is formed in the same way if step (b) yields (S)-3-chloro-2-hydroxypropio-  
5 phenone.

#### **4.2 BIOLOGICAL ACTIVITIES OF BUPROPION METABOLITES**

Bupropion metabolites can be screened for their ability to inhibit the reuptake of the neuronal monoamines norepinephrine (NE), dopamine (DA), and serotonin (5-HT).  
10 Norepinephrine reuptake inhibition can be determined using the general procedure described by Moisset, B., *et al.*, *Brain Res.*, 92:157-164 (1975); dopamine reuptake inhibition can be determined using the general procedures described by Janowsky, A., *et al.*, *J. Neurochem.* 46:1272-1276 (1986); and serotonin reuptake inhibition can be determined using the general procedures described by Perovic, S. and Muller, W. E. G.,  
15 *Brain Res.* 92:157-164 (1995).

#### **4.3. PHARMACEUTICAL COMPOSITIONS AND METHOD OF USE**

The magnitude of a prophylactic or therapeutic dose of an active ingredient in the acute or chronic management of a disorder or condition will vary with the severity of the  
20 disorder or condition to be treated and the route of administration. The dose, and perhaps the dose frequency, will also vary according to age, body weight, response, and the past medical history of the patient. Suitable dosing regimens can be readily selected by those skilled in the art with due consideration of such factors.

Suitable daily doses for the treatment or prevention of a disorder described herein  
25 can be readily determined by those skilled in the art. A recommended dose of racemic or optically pure bupropion metabolite is from about 1 mg to about 750 mg per day, given as a single once-a-day dose in the morning or as divided doses throughout the day. Preferably, a daily dose is from about 5 mg to about 700 mg per day, more preferably from about 10 mg to about 650 mg per day.

30 Suitable daily dosage ranges of second pharmacologically active compounds that can be adjunctively administered with a racemic or optically pure bupropion metabolite can

be readily determined by those skilled in the art following dosages reported in the literature and recommended in the *Physician's Desk Reference*® (53<sup>rd</sup> ed., 1999).

For example, suitable daily dosage ranges of 5-HT<sub>3</sub> antagonists can be readily determined by those skilled in the art and will vary depending on factors such as those described above and the particular 5-HT<sub>3</sub> antagonists used. In general, the total daily dose of a 5-HT<sub>3</sub> antagonist for the treatment or prevention of a disorder described herein is from about 0.5 mg to about 500 mg, preferably from about 1 mg to about 350 mg, and more preferably from about 2 mg to about 250 mg per day.

Suitable daily dosage ranges of nicotine can also be readily determined by those skilled in the art and will vary depending on factors such as those described above. In general, the total daily dose of nicotine for the treatment or prevention of a disorder described herein is from about 1 mg to about 60 mg, preferably from about 8 mg to about 40 mg, and more preferably from about 10 mg to about 25 mg per day.

The therapeutic or prophylactic administration of an active ingredient of the invention is preferably initiated at a lower dose, e.g., from about 1 mg to about 75 mg of bupropion metabolite and optionally from about 15 mg to about 60 mg of 5-HT<sub>3</sub> antagonist, and increased, if necessary, up to the recommended daily dose as either a single dose or as divided doses, depending on the global response of the patient. It is further recommended that patients aged over 65 years should receive doses of bupropion metabolite in the range of from about 1 mg to about 375 mg per day depending on global response. It may be necessary to use dosages outside these ranges, which will be readily determinable by one of ordinary skill in the pharmaceutical art.

The dosage amounts and frequencies provided above are encompassed by the terms "therapeutically effective," "prophylactically effective," and "therapeutically or prophylactically effective" as used herein. When used in connection with an amount of a racemic or optically pure bupropion metabolite, these terms further encompass an amount of racemic or optically pure bupropion metabolite that induces fewer or less severe adverse effects than are associated with the administration of racemic bupropion.

Any suitable route of administration can be employed for providing the patient with a therapeutically or prophylactically effective dose of an active ingredient. For example, oral, mucosal (e.g., nasal, sublingual, buccal, rectal, vaginal), parenteral (e.g., intravenous, intramuscular), transdermal, and subcutaneous routes can be employed. Preferred routes of

administration include oral, transdermal, and mucosal. As mentioned above, administration of an active ingredient for the treatment or prevention of erectile dysfunction is preferably mucosal or transdermal. Suitable dosage forms for such routes include, but are not limited to, transdermal patches, ophthalmic solutions, sprays, and aerosols. Transdermal

5 compositions can also take the form of creams, lotions, and/or emulsions, which can be included in an appropriate adhesive for application to the skin or can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

10 A preferred transdermal dosage form is a "reservoir type" or "matrix type" patch, which is applied to the skin and worn for a specific period of time to permit the penetration of a desired amount of active ingredient. Examples of transdermal dosage forms and methods of administration that can be used to administer the active ingredient(s) of the invention include, but are not limited to, those disclosed in U.S. Patent Nos.: 4,624,665; 4,655,767; 4,687,481; 4,797,284; 4,810,499; 4,834,978; 4,877,618; 4,880,633; 4,917,895; 15 4,927,687; 4,956,171; 5,035,894; 5,091,186; 5,163,899; 5,232,702; 5,234,690; 5,273,755; 5,273,756; 5,308,625; 5,356,632; 5,358,715; 5,372,579; 5,421,816; 5,466,465; 5,494,680; 5,505,958; 5,554,381; 5,560,922; 5,585,111; 5,656,285; 5,667,798; 5,698,217; 5,741,511; 5,747,783; 5,770,219; 5,814,599; 5,817,332; 5,833,647; 5,879,322; and 5,906,830, the disclosures of which are incorporated herein by reference.

20 An example of a transdermal dosage form of the invention comprises a bupropion metabolite and/or a second pharmacologically active compound in a patch form. The patch is worn for 24 hours and provides a total daily dose of from about 1 mg to about 750 mg per day. Preferably, a daily dose is from about 5 mg to about 700 mg per day, more preferably, from about 10 mg to about 650 mg per day. The patch can be replaced with a fresh patch. 25 when necessary to provide constant administration of the active ingredient to the patient.

Other dosage forms of the invention include, but are not limited to, tablets, coated tablets, caplets, troches, lozenges, dispersions, suspensions, suppositories, ointments, cataplasms (poultices), pastes, powders, dressings, creams, plasters, solutions, capsules, soft elastic gelatin capsules, sustained release formulations, and patches.

30 In one embodiment, pharmaceutical compositions and dosage forms of the invention comprise a racemic or optically pure bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof, and optionally a second pharmacologically active

compound, such as a SSRI, a 5-HT<sub>3</sub> antagonist, or nictotine. Preferred racemic or optically pure bupropion metabolites are (R,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (S,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (R,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; (R,R)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (S,R)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (S,S)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (R,S)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanol; (R)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanone; and (S)-1-(3-chlorophenyl)-2-[(1,1-dimethylethanol)amino]-1-propanone. The pharmaceutical compositions and dosage forms can contain a pharmaceutically acceptable carrier and optionally other therapeutic ingredients known to those skilled in the art.

In practical use, an active ingredient can be combined in an intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier can take a wide variety of forms depending on the form of preparation desired for administration. In preparing the compositions for an oral dosage form, any of the usual pharmaceutical media can be employed as carriers, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, and the like in the case of oral liquid preparations (such as suspensions, solutions, and elixirs) or aerosols; or carriers such as starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents can be used in the case of oral solid preparations, preferably without employing the use of lactose. For example, suitable carriers include powders, capsules, and tablets, with the solid oral preparations being preferred over the liquid preparations.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques.

In addition to the common dosage forms set out above, an active ingredient can also be administered by controlled release means or delivery devices that are well known to those of ordinary skill in the art, such as those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; 4,008,719; 5,674,533; 5,059,595; 5,591,767; 5,120,548;

5,073,543; 5,639,476; 5,354,556; and 5,733,566, the disclosures of which are incorporated herein by reference. These dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, or microspheres or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the pharmaceutical compositions of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include: 1) extended activity of the drug; 2) reduced dosage frequency; and 3) increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and thus can affect the occurrence of side effects.

Most controlled-release formulations are designed to initially release an amount of drug that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various inducers, including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

Pharmaceutical compositions of the invention suitable for oral administration can be presented as discrete dosage forms, such as capsules, cachets, or tablets, or aerosol sprays each containing a predetermined amount of an active ingredient as a powder or in granules, a solution, or a suspension in an aqueous or non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such dosage forms can be prepared by any of the

methods of pharmacy, but all methods include the step of bringing the active ingredient into association with the carrier, which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, 5 shaping the product into the desired presentation.

For example, a tablet can be prepared by compression or molding, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as powder or granules, optionally mixed with an excipient such as, but not limited to, a binder, a lubricant, an inert 10 diluent, and/or a surface active or dispersing agent. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

This invention further encompasses lactose-free pharmaceutical compositions and dosage forms. Because the major human metabolites of bupropion are secondary amines, 15 they can potentially decompose over time when exposed to lactose. Compositions of the invention that comprise bupropion metabolites preferably contain little, if any, lactose other mono- or di-saccharides. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

20 Lactose-free compositions of the invention can comprise excipients which are well known in the art and are listed in the USP (XXI)/NF (XVI), which is incorporated herein by reference. In general, lactose-free compositions comprise an active ingredient, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise an active ingredient, 25 microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms which comprises an active ingredient, since water can facilitate the degradation of some compounds. For example, the addition of water (*e.g.*, 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to 30 determine characteristics such as shelf-life or the stability of formulations over time. *See, e.g.*, Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, NY, 1995, pp. 379-80. In effect, water and heat accelerate decomposition. Thus the

effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

5       Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms of racemic or optically pure bupropion metabolite which contain lactose are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

10       An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastic or the like, unit dose containers, blister packs,  
15       and strip packs.

      In this regard, the invention encompasses a method of preparing a solid pharmaceutical formulation which comprises an active ingredient which method comprises admixing under anhydrous or low moisture/humidity conditions the active ingredient and an excipient (e.g., lactose), wherein the ingredients are substantially free of water. The method  
20       can further comprise packaging the anhydrous or non-hygroscopic solid formulation under low moisture conditions. By using such conditions, the risk of contact with water is reduced and the degradation of the active ingredient can be prevented or substantially reduced.

      Binders suitable for use in pharmaceutical compositions and dosage forms include,  
25       but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208,  
30       2906, 2910), microcrystalline cellulose, and mixtures thereof.

      Suitable forms of microcrystalline cellulose include, for example, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, and AVICEL-PH-105 (available

from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA, U.S.A.). An exemplary suitable binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103™ and Starch 1500 LM.

5           Examples of suitable fillers for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (*e.g.*, granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder/filler in pharmaceutical compositions of the present invention is typically present in  
10       about 50 to about 99 weight percent of the pharmaceutical composition.

          Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Too much of a disintegrant will produce tablets which may disintegrate in the bottle. Too little may be insufficient for disintegration to occur and may thus alter the rate and extent of release of the active  
15       ingredient(s) from the dosage form. Thus, a sufficient amount of disintegrant that is neither too little nor too much to detrimentally alter the release of the active ingredient(s) should be used to form the dosage forms of the compounds disclosed herein. The amount of disintegrant used varies based upon the type of formulation and mode of administration, and is readily discernible to those of ordinary skill in the art. Typically, about 0.5 to about 15  
20       weight percent of disintegrant, preferably about 1 to about 5 weight percent of disintegrant, can be used in the pharmaceutical composition.

          Disintegrants that can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin  
25       potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other algin, other celluloses, gums or mixtures thereof.

          Lubricants which can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other  
30       glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (*e.g.*, peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, or mixtures thereof. Additional lubricants include, for



example, a syloid silica gel (AEROSIL 200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, Texas), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass), or mixtures thereof. A lubricant can optionally be added, typically in an amount of less than about 1 weight percent of the pharmaceutical composition.

Dosage forms of the invention that comprise a bupropion metabolite preferably contain from about 1 mg to about 750 mg of the metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof. For example, each tablet, cachet, or capsule contains from about 1 mg to about 750 mg of the active ingredient. Most preferably, the tablet, cachet, or capsule contains either one of three dosages, *e.g.*, about 25 mg, about 50 mg, or about 75 mg of a racemic or optically pure bupropion metabolite (as scored lactose-free tablets, the preferable dose form).

The invention is further defined by reference to the following examples. It will be apparent to those skilled in the art that many modifications, both to materials and methods, can be practiced without departing from the scope of this invention.

## 5. EXAMPLES

### 5.1. EXAMPLE 1: SYNTHESIS OF (5,5)-HYDROXYBUPROPION

This synthesis, which follows that depicted in Scheme 3 of the Detailed Description, comprises the formation of three intermediates.

Z-1-(3-Chlorophenyl)-1-tert-butyldimethylsilyloxy-1-propene. A solution of LDA (33.0 mmol) in THF (100 mL) was cooled to -78°C and HMPA (5 mL) was added. The ketone [1-(3-chlorophenyl)-propanone] (8.6 g) in THF (20 mL) was slowly added over 45 minutes to this rapidly stirring mixture. After an additional 3 minutes at -78°C, TBSCl (33.0 mL, 1.0 M in hexane) was added. This mixture was stirred at -78°C for 5 minutes and allowed to warm to room temperature over 40 minutes. NaHCO<sub>3</sub> (60 mL, saturated aqueous solution) was added and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 80 mL). The organic extracts were combined, washed with brine, dried over Mg<sub>2</sub>SO<sub>4</sub> and concentrated to give a crude mixture. The product was purified by flash chromatography eluted with hexane/TEA (99.5/0.5), yielding 13.4 g product (Z/E ratio > 99). <sup>1</sup>H NMR(CDCl<sub>3</sub>): δ 0.12 (s, 6H), 0.95 (s, 9H), 2.75 (d, 3H), 5.25 (q, 1H), 7.2-7.42 (m, 4H).

- (R)-3'-Chloro-2-hydroxyl-propiofenone. Z-1-(3 -Chlorophenyl-*tert*-butyldimethylsilyloxy-1-propene (12.0 g, 44 mmol) was added to a well-stirred mixture of AD-mix-P (80 g) and  $\text{CH}_3\text{SO}_2\text{NH}_2$  (4.2 g, 44 mmol) in *tert*-butyl alcohol/water mixture (220 mL/220 mL) maintained at 0°C. The reaction mixture was stirred at 0°C for 28 hours.
- 5 Solid sodium sulfite (40 g) was added. The mixture was stirred for an additional 45 minutes and extracted with  $\text{CH}_2\text{Cl}_2$  (2 x 100 mL). The combined organic extracts were washed with  $\text{NaHCO}_3$  and brine, and evaporated. The residue was passed through a silica gel column to give the desired product (7.0 g).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.45 (d, 3H), 5.15 (q, 1H), 7.2-7.9 (m, 4H).
- 10 (S,S)-Hydroxybupropion. To a solution of (R)-3'-chloro-2-hydroxyl-propiofenone (300 mg) in  $\text{CH}_2\text{Cl}_2$  (6 mL) at -78°C was added trifluoromethanesulfonic anhydride (0.5 g), followed by addition of 2,6-lutidine (0.26 g). The reaction mixture was allowed to warm to -40°C and stirred at this temperature for 40 minutes. 2-Amine-2-methyl-1-propanol (0.4 g, 2.5 eq) was added, and stirred for 2 hours at -40°C. The reaction mixture was warmed to
- 15 room temperature and stirred overnight. It was extracted with  $\text{CH}_2\text{Cl}_2$  (10 mL). The extract was washed with  $\text{NaHCO}_3$ , water, and brine, concentrated to give a residue. The final product was purified by chromatography eluted with  $\text{CH}_3\text{CN}$  (180 mg, e.e. > 99%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.78 (d, 3H), 1.1 (s, 3H), 1.4 (s, 3H), 3.2 (q, 1H), 3.4 (d, 1H), 3.8 (d, 2H), 7.2-7.65 (m, 4H).  $[\alpha] = +66^\circ$  (c = 1, EtOH). (S,S)-hydroxybupropion free base was treated
- 20 with HCl in diethyl ether to give its HCl salt.  $[\alpha] = \pm 30.6^\circ$  (c = 1, EtOH).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  1.0 (d, 3H), 1.32 (s, 3H), 1.56 (s, 3H), 3.4 (s, 1H), 3.4 (d, 1H), 4.0 (d, 1H), 7.5 (m, 5H), 8.8 (s, 1H), 10.1 (s, 1H). e.e. 99.4% as determined by HPLC with chiral column, ChiralCEL GD. 4.6 x 250 mm, 10 nm, hexane/ethanol/diethylamine (98:2:0.1). (R,R)-hydroxybupropion was prepared from (S)-3'-chloro-2-hydroxyl-
- 25 propiofenone with 97% e.e. as determined by HPLC with chiral column, ChiralCEL GD. 4.6 x 250 mm, 10 nm, hexane/ethanol/diethylamine (98:2:0.1).

## 5.2. EXAMPLE 2: NEURONAL MONOAMINE REUPTAKE INHIBITION

- The abilities of racemic bupropion [BP( $\pm$ )], and the bupropion metabolites (S,S)-
- 30 hydroxybupropion [HBP(S,S)], (R,S)-hydroxybupropion [HBP(R,S)], and (RS,RS)-hydroxybupropion [HBP(RS,RS)] to inhibit the reuptake of neuronal monoamines was determined using the general methods of Moisset, B., *et al.*, *Brain Res.* 92:157-164 (1975),

Janowsky, A., *et al.*, *J. Neurochem.* 46:1272-1276 (1986), and Perovic, S. and Muller, W. E.G., *Brain Res.* 92:157-164 (1995).

Inhibition of norepinephrine (NE) reuptake was determined using rat hypothalamus as a tissue source and protryptiline as a reference compound. Inhibition of dopamine (DA) reuptake was determined using rat corpora striata as a tissue source and GBR 12909 as a reference compound. Inhibition of serotonin (5-HT) reuptake was determined using rat brain as a tissue source and imipramine as a reference compound. The specific conditions for each assay are shown in Table 1:

10

Table 1

Assay	Substrate	Incubation	Detection Method
NE	[ <sup>3</sup> H]NE (0.2 $\mu$ Ci/mL)	20 min. / 37°C	liquid scintillation
DA	[ <sup>3</sup> H]DA (0.2 $\mu$ Ci/mL)	15 min. / 37°C	liquid scintillation
5-HT	[ <sup>3</sup> H]5-HT (0.2 $\mu$ Ci/mL)	15 min. / 37°C	liquid scintillation

15

wherein the end products observed were formed by the incorporation of [<sup>3</sup>H]NE, [<sup>3</sup>H]DA, and [<sup>3</sup>H]5-HT into synaptosomes. Radioactivity was determined with a scintillation counter (Topcount, Packard) using a liquid scintillation cocktail (Microscint 0, Packard).

Racemic bupropion and the bupropion metabolites were first tested in each assay at 10  $\mu$ M in duplicate or triplicate. For assays wherein they inhibited the reuptake by more than 50% at this concentration, they were further tested at eight concentrations in duplicate to obtain full inhibition curves. In each experiment, the respective reference compound was tested at eight concentrations in duplicate to obtain an inhibition curve in order to validate this experiment.

IC<sub>50</sub> values and Hill coefficients (nH) were determined for the reference compounds and the test compounds (*i.e.*, bupropion and metabolites of bupropion) by non-linear regression analysis of their inhibition curves. These parameters were obtained by Hill equation curve fitting.

None of the compound tested significantly inhibited 5-HT reuptake. The IC<sub>50</sub> values determined for these compounds with regard to norepinephrine and dopamine reuptake are presented in Table 2:

30

Table 2

Compounds	NE reuptake		DA reuptake	
	IC <sub>50</sub> (nM)	(nH)	IC <sub>50</sub> (nM)	(nH)
HBP(S,S)	229	(0.8)	1,400	(1.0)
BP(±)	756	(1.1)		
HBP(R,S)	746	(>3)	294	(0.9)
HBP(R,R)	-		-	
	IC <sub>50</sub> (nM)	(nH)	IC <sub>50</sub> (nM)	(nH)
protriptyline	3.6/3.8	(2.6)/(1.4)		
GBR 12909			5.6	(1.7)

10

The measured biological activity of the bupropion metabolites are unexpectedly different than the activity of bupropion itself. For example, racemic bupropion (*i.e.*, (±)1-(3-chlorophenyl)-2-[(1,1-dimethylethyl)amino]-1-propanone) inhibits norepinephrine reuptake with an IC<sub>50</sub> of approximately 746 nM, while the optically pure metabolite (S,S)-hydroxybupropion (*i.e.*, (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol) inhibits norepinephrine with a dramatically lower IC<sub>50</sub> of 229 nM. And while racemic bupropion inhibits dopamine reuptake with an IC<sub>50</sub> of approximately 294 nM, the optically pure metabolite (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol does not significantly inhibit dopamine reuptake, having an IC<sub>50</sub> of approximately 1400 nM. But like racemic bupropion, this optically pure metabolite does not measurably inhibit serotonin reuptake.

These results indicate that the biological activity of each of the bupropion metabolites of the invention is dramatically and unexpectedly different from that of bupropion. These results further indicate that bupropion metabolites are superior in their abilities to treat certain disorders. For example, optically pure (S,S)-hydroxybupropion is surprisingly selective with regard to its inhibition of neuronal monoamine reuptake, and may thus be used to inhibit norepinephrine reuptake.

### 5.3. EXAMPLE 3: *IN VIVO* ACTIVITY: SEIZURE MODEL

The pharmacological effects of a bupropion metabolite can be determined in a number of ways. For example, its ability to inhibit artificially induced seizures in mice may be informative.

- 5        Using the methods of Green and Murray, *J. Pharm. Pharmacol.* 41:879-880 (1989), a group of 4-6 rats are lightly restrained and a 10 mg/mL solution of the convulsant drug pentetrazol is infused via a 25 gauge needle inserted into a tail vein of each rat at a rate of 2.6 mL/min. The time of infusion of the convulsant drug required to produce the first myoclonic twitch (which occurs with the first EEG abnormality) is recorded and doses  
10       required to produce the seizure calculated. Seizure threshold is expressed as mg/kg and can be calculated using the formula:

$$(I \times C \times T) / (60 \times W)$$

wherein I is the infusion rate measured in mL per minute; C is the drug concentration in 10 mg/mL; T is the time to twitch in seconds; and W is the rate weight in kilograms.

- 15       Bupropion metabolites are administered by intraperitoneal or intravenous injection 15 minutes before the determination of seizure threshold.

### 5.4. EXAMPLE 4: *IN VIVO* ACTIVITY: PHENYLOQUINONE WRITHING ASSAY

- 20       The pharmacological effects of a bupropion metabolite can also be determined from the antiphenylquinone writhing test, which is a standard procedure for detecting and comparing analgesic activity in laboratory animals. The advantage of this test is that it generally correlates well with human efficacy. In response to an injected, locally irritating  
25       solution, the animals have writhings that are inhibited by analgesic agents.

- Mice dosed with at least two dose levels of a bupropion metabolite are challenged with phenyl-p-benzoquinone (PPQ) given intraperitoneally and then observed for the characteristic stretch-writhing syndrome. Lack of writhing constitutes a positive response. The degree of analgesic protection can be calculated on the basis of suppression of writhing  
30       relative to control animals run on the same day. Time response data are also obtained. Observations are made early enough after dosing to detect differences in onset.

For example, the following protocols may be used, wherein ten mice are used per dose group:

Preparation of Phenylquinone: PPQ is made-up as a 0.02% aqueous solution in ethyl alcohol. PPQ (20 mg) is ground and dissolved in a tissue homogenizer in 5 mL ethyl alcohol, and the volume brought to 100 mL with distilled water, preheated to 45°C. The resulting solution should be a clear amber color. PPQ solutions are made fresh twice daily and, if necessary, about every four hours because of the tendency of PPQ to precipitate out of solution.

Dose amounts: 0.1, 0.3, 1.0, 3.0, 10.0, 30.0, and 100.0 mg/kg.

Positive Control: Aspirin - 200 mg/kg.

Writhing: PPQ solution is administered intraperitoneally using a 25 gauge, 5/8" long needle on a 1 mL syringe. Each animal in the group receives 0.25 mL. The group of ten mice per dose level is observed closely for ten minutes for exhibition of writhing. The stability of the PPQ solution(s) to produce the writhing response is verified for each preparation in ten mice to which the vehicle was administered prior to PPQ administration.

Characteristic patterns of writhing consist of torsion of the abdomen and thorax, drawing the hind legs close to the body and raising the heels of the hind feet off of the floor.

Observation Times: Reference and positive control article activity is studied at 60 minutes after administration. After the designated absorption time interval of a group has elapsed, the mice are challenged with PPQ. Each mouse receives one dose of 0.25 mL of PPQ. After PPQ administration, the mouse is placed in individual Plexiglas® squares 4" x 4" x 5" deep and observed closely for a ten minute period for exhibition of the writhing syndrome.

Scoring Determinations: The total number of writhes for each mouse is recorded. The mean number of writhes for the control and each positive control and reference group is compared and percent inhibition calculated.

25

### 5.5. EXAMPLE 5: IN VIVO ACTIVITY: FORMALIN TEST

The pharmacological effects of a bupropion metabolite may also be determined from other models, some of which are discussed by Bannon, A.W., *et al.*, *Science* 279:77-81 (1998). One of these models is the formalin test.

The formalin test is an animal model for persistent inflammatory pain. In the formalin test, the second phase of the biphasic nociceptive response is thought to be

mediated in part by a sensitization of neuronal function at the level of the spinal cord, and reflect the clinical observation of hyperalgesia associated with tissue injury.

Using the method of Dubusson, D., and Dennis, S.G., *Science* 4:161 (1977), rats are allowed to acclimate to their individual cages for 20 minutes, after which time 50 mL of a 5% formalin solution is injected into the dorsal aspect of one of the rear paws. The rats are then returned to the clear observation cages, which are suspended above mirror panels. Only phase 2 of the formalin test may be scored, and phase 2 may be defined as the 20-minute period of time from 30 to 50 minutes after formalin injection. The investigator records nocifensive behaviors in the injected paw of four animals during the session by observing each animal for one 15-second observation period during each 1-minute interval. Nocifensive behaviors include flinching, licking, or biting the injected paw. In dose-response studies, the test compound (or saline) is administered 5 minutes before the injection of formalin. In antagonist studies, the antagonists or saline are administered 10 minutes before treatment.

#### 5.6. EXAMPLE 6: *IN VIVO* ACTIVITY: NEUROPATHIC PAIN MODEL

Another pharmacological model discussed by Bannon, A.W., *et al.*, *Science* 279:77-81(1998) is the neuropathic pain test. In the neuropathic pain model, nerve injury results in neuroplastic changes that lead to allodynia, a condition characterized by nocifensive behavioral responses to what are normally non-noxious stimuli conducted by A $\beta$  fibers. In the Chung model of neuropathic pain, allodynia is produced in the hind limb ipsilateral by ligation of the L5 and L6 spinal nerves. S.H. Kim and J.M. Chung, *Science* 50, 355 (1992). According to this model, a within-subjects design in which all animals receive all treatments is used for dose-response studies.

Using the Chung model, baseline allodynia scores are determined for all animals before the start of the drug studies. Only rats with threshold scores are considered allodynic and used in further testing. Drug studies (separate studies for each compound) begin approximately 2 weeks after nerve ligation surgery. For dose-response experiments, animals are tested over a 2-week period. Test days are separated by 2 to 3 day intervals during which no testing is conducted and no treatment is given. On test days, animals are placed in individual chambers and allowed to acclimate for 15 to 20 minutes. After acclimation, baseline scores are determined. Next, animals are treated and scores are

determined 15, 30, 50, and 120 minutes after treatment. This procedure is repeated on test days until each animal has received all treatments for any given drug. The treatment order is counterbalanced across animals. For statistical analysis, the time point of peak effect is compared.

5

### 5.7. EXAMPLE 7: ORAL FORMULATION

Table 3 provides the ingredients for a lactose-free tablet dosage form of a bupropion metabolite:

Table 3

10	Component	Quantity per Tablet (mg)
	Bupropion metabolite (e.g., (S,S)-hydroxybupropion)	75
	Microcrystalline cellulose	125
	Talc	5.0
15	Water (per thousand tablets)	30.0 mL *
	Magnesium Stearate	0.5

\* The water evaporates during manufacture.

The active ingredient (bupropion metabolite) is blended with the cellulose until a uniform blend is formed. The smaller quantity of corn starch is blended with a suitable quantity of water to form a corn starch paste. This is then mixed with the uniform blend until a uniform wet mass is formed. The remaining corn starch is added to the resulting wet mass and mixed until uniform granules are obtained. The granules are then screened through a suitable milling machine, using a 1/4 inch stainless steel screen. The milled granules are then dried in a suitable drying oven until the desired moisture content is obtained. The dried granules are then milled through a suitable milling machine using 1/4 mesh stainless steel screen. The magnesium stearate is then blended and the resulting mixture is compressed into tablets of desired shape, thickness, hardness and disintegration. Tablets are coated by standard aqueous or nonaqueous techniques.

Another tablet dosage formulation suitable for use with the active ingredients of the invention is provided by Table 4:



Table 4

Component	Quantity per Tablet (mg)		
	Formula A	Formula B	Formula C
Bupropion metabolite ( <i>e.g.</i> , (S,S)-hydroxybupropion)	20	40	100
Microcrystalline cellulose	134.5	114.5	309.0
Starch BP	30	30	60
Pregelatinized Maize Starch BP	15	15	30
Magnesium Stearate	0.5	0.5	1.0
Compression Weight	200	200	500

The active ingredient is sieved and blended with cellulose, starch, and pregelatinized maize starch. Suitable volumes of purified water are added and the powders are granulated. After drying, the granules are screened and blended with the magnesium stearate. The granules are then compressed into tablets using punches.

Tablets of other strengths may be prepared by altering the ratio of active ingredient to pharmaceutically acceptable carrier, the compression weight, or by using different punches.

#### 5.8. EXAMPLE 8: ORAL FORMULATION

Table 5 provides the ingredients for a capsule dosage form of a bupropion metabolite:

Table 5

Component	Quantity per Tablet (mg)		
	Formula A	Formula B	Formula C
Bupropion metabolite (e.g., (S,S)-hydroxybupropion)	25	50	75
Microcrystalline cellulose	149.5	124.5	374
Corn Starch	25	25	50
Water (per thousand tablets)	0.5	0.5	1.0
Magnesium Stearate	200	200	200

10           The active ingredient, cellulose, and corn starch are blended until uniform; then the magnesium stearate is blended into the resulting powder. The resulting mixture is encapsulated into suitably sized two-piece hard gelatin capsules using suitable machinery. Other doses can be prepared by altering the ratio of active ingredient to pharmaceutically acceptable carrier, the fill weight, and, if necessary, by changing the capsule size to suit.

15           The active ingredient, cellulose, and corn starch are blended until uniform; then the magnesium stearate is blended into the resulting powder. The resulting mixture is encapsulated into suitably sized two-piece hard gelatin capsules using suitable machinery. Other doses can be prepared by altering the ratio of active ingredient to pharmaceutically acceptable carrier, the fill weight, and, if necessary, by changing the capsule size to suit.

20           The embodiments of the invention described above are intended to be merely exemplary and those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein. All such equivalents are considered to be within the scope of the invention and are encompassed by the following claims.

25

## THE CLAIMS

What is claimed is:

- 5           1.     A method of treating or preventing a disorder that is ameliorated by the inhibition of neuronal monoamine reuptake which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite, or a pharmaceutically acceptable salt, solvate, or clathrate thereof.
- 10           2.     The method of claim 1 wherein the bupropion metabolite is optically pure.
3.     The method of claim 2 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.
- 15           4.     The method of claim 1 wherein the adverse effects associated with the inhibition of dopamine reuptake are reduced or avoided.
5.     The method of claim 1 wherein the bupropion metabolite or
- 20     pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively administered with a second pharmacologically active compound.
6.     A method of treating or preventing erectile dysfunction which comprises administering to a patient in need of such treatment or prevention a therapeutically or
- 25     prophylactically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.
7.     The method of claim 6 wherein the bupropion metabolite is optically pure.
- 30           8.     The method of claim 7 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.

9. The method of claim 6 wherein the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered transdermally or mucosally.

5 10. The method of claim 6 wherein the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively administered with a 5-HT<sub>3</sub> antagonist.

10 11. The method of claim 10 wherein the 5-HT<sub>3</sub> antagonist is an antiemetic agent.

12. The method of claim 10 wherein the 5-HT<sub>3</sub> antagonist is selected from the group consisting of granisetron, metoclopramide, ondansetron, renzapride, zacopride, tropisetron, and optically pure stereoisomers, active metabolites, and pharmaceutically acceptable salts, solvates, and clathrates thereof.

15 13. A method of treating or preventing an affective disorder which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.

20 14. The method of claim 13 wherein the bupropion metabolite is optically pure.

15. The method of claim 14 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.

25 16. The method of claim 13 wherein the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively administered with a therapeutically or prophylactically effective amount of a second pharmacologically active compound.

30 17. The method of claim 13 wherein the affective disorder is depression.

18. The method of claim 13 wherein the affective disorder is narcolepsy.
19. The method of claim 13 wherein the affective disorder is nicotine addiction.
- 5 20. A method of treating or preventing a cerebral function disorder which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.
- 10 21. The method of claim 20 wherein the bupropion metabolite is optically pure.
22. The method of claim 21 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.
- 15 23. The method of claim 20 wherein the the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively administered with a therapeutically or prophylactically effective amount of a second pharmacologically active compound.
- 20 24. The method of claim 20 wherein the cerebral function disorder is Parkinson's disease.
25. The method of claim 20 wherein the cerebral function disorder is epilepsy.
- 25 26. A method of eliciting smoking cessation which comprises administering to a patient who smokes tobacco a therapeutically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.
27. The method of claim 26 wherein the bupropion metabolite is optically pure.
- 30 28. The method of claim 27 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.

29. The method of claim 26 wherein the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered orally, mucosally, or transdermally.

5           30. The method of claim 29 wherein the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered transdermally.

31. The method of claim 26 wherein the bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is adjunctively administered  
10 with a therapeutically effective amount of nicotine.

32. The method of claim 31 wherein the nicotine and/or bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered orally, mucosally, or transdermally.

15           33. The method of claim 32 wherein the nicotine and/or bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof is administered transdermally.

34. A method of treating or preventing incontinence which comprises  
20 administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.

35. The method of claim 34 wherein the bupropion metabolite is optically pure.

25           36. The method of claim 35 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.

37. The method of claim 34 wherein incontinence is stress urinary incontinence.

30           38. The method of claim 34 wherein the patient is a human of an age greater than 50 years or less than 13 years.

39. A pharmaceutical composition which comprises a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.

40. The pharmaceutical composition of claim 39 wherein the bupropion  
5 metabolite is optically pure.

41. The pharmaceutical composition of claim 40 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.

10 42. The pharmaceutical composition of claim 40 wherein said pharmaceutical composition further comprises a second pharmacologically active compound selected from the group consisting of selective serotonin reuptake inhibitors, 5-HT<sub>3</sub> inhibitors, and nicotine.

15 43. A dosage form comprising a bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.

44. The dosage form of claim 43 wherein said dosage form is solid.

20 45. The dosage form of claim 43 wherein said dosage form is a sterile solution.

46. The dosage form of claim 43 wherein said dosage form is a transdermal patch.

25 47. The dosage form of claim 43 wherein the bupropion metabolite is optically pure.

48. The dosage form of claim 47 wherein the optically pure bupropion metabolite is optically pure (S,S)-hydroxybupropion.

30

49. The dosage form of claim 43 wherein said dosage form further comprises a second pharmacologically active compound selected from the group consisting of selective serotonin reuptake inhibitors, 5-HT<sub>3</sub> inhibitors, and nicotine.

5 50. The dosage form of claim 43 wherein said dosage form is suitable for oral, mucosal, or transdermal administration to a patient.

10 51. A dosage form suitable for transdermal administration to a patient which comprises nicotine and a bupropion metabolite or pharmaceutically acceptable salt, solvate, or clathrate thereof.

52. A lactose-free solid dosage form comprising an optically pure bupropion metabolite or a pharmaceutically acceptable salt, solvate, or clathrate thereof.

15 53. The dosage form of claim 52 wherein said dosage form is an oral dosage form.

20 54. A process for preparing optically pure (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol or a pharmaceutically acceptable salt, solvate or clathrate thereof which comprises:

asymmetrically hydroxylating Z-1-(3-chlorophenyl)-1-*tert*-butyldimethylsilyloxy-1-propene to form an intermediate;

reacting the intermediate with 2-amino-2-methyl-1-propanol to form (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; and

25 isolating the (S,S)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol.



55. A process for preparing optically pure (R,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol or a pharmaceutically acceptable salt, solvate or clathrate thereof which comprises:


asymmetrically hydroxylating Z-1-(3-chlorophenyl)-1-*tert*-butyldimethylsilyloxy-1-propene to form an intermediate;

reacting the intermediate with 2-amino-2-methyl-1-propanol to form (R,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol; and

isolating the (R,R)-2-(3-chlorophenyl)-2-hydroxy-3,5,5-trimethyl-morpholinol.

56. The process of claim 54 or 55 wherein the intermediate formed by the asymmetric hydroxylation of Z-1-(3-chlorophenyl)-1-*tert*-butyldimethylsilyloxy-1-propene is an  $\alpha$ -hydroxy ketone activated by trifluoromethane sulfonic anhydride.

57. A compound of the formula:



20

wherein R is selected from the group consisting of hydrogen, triflate, tosylate, and nosylate; or a pharmaceutically acceptable salt, solvate, or clathrate thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 September 2000 (08.09.2000)

PCT

(10) International Publication Number  
**WO 00/51546 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 31/5375**,  
31/133, 31/137, C07C 49/76, 49/82, 49/807, 317/14,  
C07D 265/32

(21) International Application Number: **PCT/US00/05109**

(22) International Filing Date: 29 February 2000 (29.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/122,277 1 March 1999 (01.03.1999) US  
60/148,324 11 August 1999 (11.08.1999) US  
Not furnished 22 February 2000 (22.02.2000) US

(71) Applicant: **SEPRACOR INC.** [US/US]; 111 Locke  
Drive, Marlborough, MA 01752 (US).

(72) Inventors: **JERUSSI, Thomas, P.**; 19 Garvey Road,  
Framingham, MA 01701 (US). **MCCULLOUGH, John,**  
**R.**; 6 Davidson Road, Worcester, MA 01605 (US).  
**SENANAYAKE, Chrisantha, H.**; 11 Old Farm Circle,  
Shrewsbury, MA 01545 (US). **FANG, Qun, K.**; 35 At-  
wood Street, Wellesley, MA 02181 (US).

(74) Agents: **INSOGNA, Anthony, M. et al.**; Pennie & Ed-  
monds LLP, 1155 Avenue of the Americas, New York, NY  
10036 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,  
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,  
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,  
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,  
UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments.

(88) Date of publication of the international search report:  
11 January 2001

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

**WO 00/51546 A3**

(54) Title: **BUPROPION METABOLITES AND METHODS OF THEIR SYNTHESIS AND USE**

(57) Abstract: Methods and compositions are disclosed which utilize metabolites of bupropion for treating disorders ameliorated by inhibition of neuronal monoamine reuptake. Such disorders include, but are not limited to, erectile dysfunction, affective disorders, cerebral function disorders, cigarette smoking, and incontinence. The invention further discloses methods of making optically pure bupropion metabolites.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/05109

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/5375, 133, 137; C07C 49/76, 82, 807; 317/14; C07D 265/32

US CL : 514/239.5, 653, 649; 544/173; 568/31, 336

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/239.5, 653, 649; 544/173; 568/31, 336

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MUSSO et al. Synthesis and Evaluation of the Anticonvulsant Activity of a Series of 2-Amino-1-Phenyl-1-Propanols Derived from the Metabolites of the Antidepressant Bupropion. Bioorganic & Medicinal Chemistry Letters. 07 January 1997, Vol. 7, No. 1, pages 1-6, see entire document.	1-5, 20-25, 39-56
Y --- A	KELLY et al. (2S 35, 5K)-2-(3,5-Difluorophenyl)-3,5-dimethyl-2-morpholinol: A Novel Antidepressant Agent and Selective Inhibitor of Norepinephrine Uptake. J. Med. Chem. 19 January 1996, Vol. 39, No. 2, pages 347-349, see entire document.	1-5, 13-19, 39-56 ----- 6-12, 20-38, 57



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents.	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 MAY 2000

Date of mailing of the international search report

02 NOV 2000

Name and mailing address of the ISA/US...  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PHYLLIS SPIVACK

Telephone No. (703) 308-1233